

Clinical profile of dengue infection at a teaching hospital in North India

Ritu Karoli, Jalees Fatima, Zeba Siddiqi, Khursheed I. Kazmi, Amit R. Sultania

Department of Medicine, Era's Lucknow Medical College, Sarfarazganj, Hardoi Road, Lucknow, Uttar Pradesh, India

Abstract

Introduction: Dengue viral infections are among the most important mosquito-borne diseases of the Indian subcontinent and have become a major global public health concern. Spread of disease has led to increased recognition of atypical manifestations apart from the classical clinical features of dengue infection.

Methodology: A cross-sectional study of admitted patients suspected to have dengue infection was conducted during the monsoon and post-monsoon seasons in the year 2010. Patients who had serological confirmation of dengue infection were classified according to World Health Organization definitions of dengue fever and dengue hemorrhagic fever. Clinical and biochemical parameters were compared between the two groups.

Results: Out of 356 patients with suspected dengue fever enrolled in the study, 138 (39%) had serologically confirmed dengue infection. Eighty (58%) patients were males and 58 (42%) were females. Ninety-six (70%) patients had classical dengue fever while 42 (30%) had dengue hemorrhagic fever. The most common symptoms were headache (105, 76%), abdominal pain (87, 63%), vomiting (80, 58%), rash (36, 26%), and cutaneous hypersensitivity (22, 16%). Hemorrhagic manifestations were present in 55 (40%) patients. Atypical manifestations were recorded. Notably, 14% of patients had neurological involvement and 4% had acute hepatic failure. Overall mortality was 6% and all fatal cases were due to multi-organ failure.

Conclusion: Dengue infection poses a huge burden to the health-care system; its spectrum ranges from mild self-limiting illness to severe fatal disease. It can have varied and multi-systemic manifestations which can go unrecognized. Clinicians should have a high index of suspicion for atypical manifestations.

Key words: dengue infection; dengue hemorrhagic fever; atypical manifestations of dengue fever; hepatic failure in dengue

J Infect Dev Ctries 2012; 6(7):551-554.

(Received 01 April 2011-- Accepted 26 July 2011)

Copyright © 2012 Karoli *et al.* This is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

In recent years, dengue has become a major global public health concern. Approximately 2.5 billion people, living mainly in urban areas of tropical and subtropical regions, are estimated to be at risk of acquiring dengue infection [1]. While dengue is endemic in more than 100 countries, most cases are reported from Southeast Asia and the western Pacific regions [2]. The resurgence of dengue has been observed in India and dengue outbreaks have been frequently reported from different parts of the country in both urban and rural populations [3-8]. Dengue infections vary in severity, ranging from influenza-like self-limiting illness to life-threatening dengue hemorrhagic fever (DHF) and dengue shock syndrome (DSS) which, if left untreated, are associated with mortality as high as 20% [2]. The various manifestations of dengue may not have a distinct line of demarcation: apart from the classic features, reports of rare presentations have

recently become more frequent [9,10]. Some presentations that are not classifiable under the World Health Organization (WHO) definitions may be potentially serious and may lead to increased morbidity and mortality of the disease. Many of these manifestations may remain unrecognized and unreported due to lack of awareness among primary care physicians. The most recent dengue outbreak was observed in the northern part of the country last year during the monsoon and post-monsoon period. We undertook this prospective study in the Department of Medicine, Era's Lucknow Medical College, from July to October 2010 to assess the clinical profile of dengue infection in hospitalized patients as well as to observe rare manifestations of dengue fever in the current outbreak and compare them with the WHO dengue classification [11].

Table 1. Comparison of clinical parameters between dengue fever and dengue hemorrhagic fever

	Total dengue positive (n = 138)	Dengue fever (n = 96)	Dengue hemorrhagic fever (n = 42)	P value
Headache/Myalgia	105 (76%)	77 (80%)	28 (66%)	0.57
Hemorrhagic manifestations	55 (40%)	13 (14%)	42 (100%)	0.01*
Rash	38 (28%)	26 (27%)	12 (29%)	0.82
Hepatomegaly	72 (52%)	47 (49%)	25 (60%)	0.52
Signs of plasma leakage	42 (30%)	-	42 (100%)	-

*p < 0.05

Methodology

We included all suspected dengue patients of 14 years or older admitted to medical wards and who appeared to have acute febrile illness with myalgia, arthralgia, headache, retro orbital pain, abdominal pain, nausea and vomiting, bleeding, hypotension or thrombocytopenia. A detailed history as well as a general and systemic clinical examination (including the tourniquet test) was recorded. Haematological profiles and biochemical investigations were done at the time of admission and were followed by daily (or bi-daily) investigations as required until discharge. Signs of plasma leakage were assessed by chest radiograph and abdominal ultrasonography. Specific investigations were performed in patients who presented with neurological involvement (cerebrospinal fluid analysis, neuroimaging, electrodiagnostic studies or muscle biopsy) or hepatic failure (viral markers, peripheral smear and serology for plasmodium falciparum, typhoid fever and leptospirosis).

Patients were classified as dengue fever, dengue hemorrhagic fever or dengue shock syndrome according to WHO guidelines [11] and laboratory diagnosis of dengue was established by demonstration of specific IgM antibodies to dengue in serum.

The study enrolled 356 patients of suspected dengue fever of whom 138 (39%) were serologically confirmed to have dengue infection. Eighty (58%) patients were males and 58 (42%) were females. Ninety-six (70%) patients had classic dengue fever while 42 (30%) fulfilled the criteria of dengue hemorrhagic fever. Of those patients with dengue hemorrhagic fever, eight patients had developed dengue shock syndrome.

Statistical analysis

Statistical analysis was performed by Chi Square test done by using the Statistical Package for Social

Sciences (SPSS 15.0 version, IBM, Chicago, USA) with $p < 0.05$ taken as statistically significant.

Ethical considerations

The study was approved by the Institutional Ethics Committee and written informed consent was obtained from all the patients

Results

Clinical diagnosis of dengue hemorrhagic fever was based on the presence of the following four criteria: (i) continuous high-grade fever lasting 2 to 7 days; (ii) hemorrhagic tendency as shown by a positive tourniquet test, petechiae or epistaxis; (iii) thrombocytopenia $< 100,000/\mu\text{l}$; (iv) evidence of plasma leakage manifested by hemoconcentration (an increase in hematocrit 20% above the average for age, sex and population); pleural effusion and ascites, *etc.* Severity was graded from mildest to most severe depending on the basis of the presence of hypotension or circulatory failure, the presence of cold clammy skin and restlessness, tachycardia and weak pulse with pulse pressure $< 20\text{mmHg}$ from mildest to severe degree. Dengue shock syndrome was diagnosed if there was profound shock and pulse and blood pressure were not recordable.

In our study the mortality rate was 6 % (8) and all fatal cases were due to multi-organ failure. The most common symptoms were headache (105, 76%), abdominal pain (87, 63%), vomiting (80, 58%), rash (36, 26%), and cutaneous hypersensitivity (22, 16%). Haemorrhagic manifestations were present in 55 (40%) patients in which epistaxis was the most common symptom. Per laboratory parameters, 119 (86%) patients had leucopenia; 123 (89%) had thrombocytopenia; 127 (92%) had elevation of liver enzymes; 66 (48%) had raised serum bilirubin; an altered coagulation profile was found in 47 (34%) patients; and 4% (6) patients developed acute fulminant hepatic failure. The clinical and

Table 2. Comparison of biochemical parameters between dengue fever and dengue hemorrhagic fever

	Total dengue positive (n = 138)	Dengue fever (n = 96)	Dengue hemorrhagic fever (n =42)	P value
Thrombocytopenia(platelet count <100,000/ μ l	123 (89%)	81 (84%)	42 (100%)	0.52
Abnormal PT/APTT	47 (34%)	13 (14%)	34 (81%)	0.01*
Elevation of transaminases	127 ((92%)	87 (90%)	40 (95%)	0.87
Hypoproteinemia (<5.5gm/dl)	47 (34%)	12 (13%)	35 (83%)	0.01*

PT-Prothrombin time APTT-Activated partial thromboplastin time

*p < 0.05

biochemical parameters in patients with dengue fever and dengue hemorrhagic fever are shown in Tables 1 and 2.

Apart from the classical manifestations of the dengue infection, we particularly observed certain rare and atypical manifestations in this outbreak. Neurological complications were found in 14% of our patients compared to the previously reported 0.5-6% [12]. Neurological involvement was present in the form of encephalopathy (6), hypokalaemic periodic paralysis (2), myositis (1) and Gullain-Barre syndrome (1). This neurological involvement can be related to the neurotrophic effect of the virus, the systemic effects of dengue infection, or the host immune response [13].

Discussion

Liver enzyme elevation, a common feature in dengue infection [14,15] was also apparent in our study. In this study, AST levels were equal to or greater than those of ALT levels in all of dengue infected patients, a finding that has also been reported earlier [16]. Acute hepatic failure, a rarely reported manifestation of dengue hemorrhagic fever [17], was diagnosed in six of our patients. Deranged liver function in dengue infection can be a result of the direct effect of the virus on liver cells or the unregulated host immune response against the virus. Fulminant hepatic failure occurs because of acute severe hepatitis and massive necrosis of the liver, causing hepatic encephalopathy and even death.

A significant proportion of patients (84%) with classical or uncomplicated dengue fever had thrombocytopenia and 13% of them had an altered coagulation profile as well. Overall, an altered coagulation profile was observed in 34% patients in our study and is indicative of the activation of both coagulation and fibrinolysis during acute dengue

infection, which is found to be particularly greater in patients with dengue hemorrhagic fever [18].

The application of WHO the classification system is not as simple and straightforward as it seems and clinical features may overlap among different categories. The WHO classification system of dengue does not include unusual manifestations such as encephalopathy, acute hepatic failure, cardiomyopathy and acute respiratory distress syndrome, which might be life-threatening. Although these manifestations are rare, they have been reported from endemic regions [9,19]. Therefore, clinicians should have a high index of suspicion and knowledge of these atypical manifestations, particularly in view of the increasing burden of dengue on the health-care system.

References

- Halstead SB (2007) Dengue. *Lancet* 370: 1644-1652.
- WHO (2009) Dengue Guidelines for Diagnosis, Treatment, Prevention and Control WHO (2009) http://whqlibdoc.who.int/publications/2009/9789241547871_eng.pdf. Last accessed 5 July 2012.
- Kabra SK, Jain Y, Pandey RM, Madhulika, Singhal T, Tripathi P, Broor S, Seth P, Seth V (1999) Dengue hemorrhagic fever in children in the 1996 Delhi epidemic. *Trans R Soc Trop Med Hyg* 73: 435-440.
- Narayan M, Aravind MA, Thilothammal N, Prema R, Sargunam CS, Ramamurty N (2002) Dengue fever epidemic in Chennai: a clinical profile and outcome. *Indian Pediatr* 39: 1027-1033.
- Shah I, Deshpande GC, Tardeja PN (2002). Outbreak of dengue in Mumbai and predictive markers for dengue shock syndrome. *J Trop Pediatr* 50: 301-305.
- Gupta P, Khare V, Tripathi S, Nag VL, Kumar R, Khan MY, Dhole TK (2010) Assessment of WHO definition of dengue hemorrhagic fever in North India. *J Infect Dev Ctries* 4: 150-155.
- Kumar A, Sharma SK, Padbidri VS, Thakrey JP, Jain DC and Dutta KK (2001) An outbreak of dengue fever in rural areas of northern India. *J Commun Dis* 233: 274-281.
- Arunachalam N, Murty US, Kabilan L, Balasubramaniyam A, Thenmozhi V, Narhari D, Ravi A, Satyanarayana K (2004) Studies on dengue in rural areas of Kurnool District,

- Andhra Pradesh, India. *J Am Mosq Control Assoc* 20: 87-90.
9. Gulati S and Maheshwari A (2007) Atypical manifestations of dengue. *Trop Med Int Health* 12: 1087-1095.
 10. Misra UK, Kalita J, Syam UK, Dhole TN (2006) Neurological manifestations of dengue virus infection. *J Neurol Sci* 244: 117-122.
 11. World Health Organization, Geneva (1997) Dengue hemorrhagic fever: Diagnosis, treatment prevention and control. 2nd edition: 12-23.
 12. Hendarto SK and Hadinegoro SR (1992) Dengue encephalopathy. *Acta Paediatr Jpn* 34: 350-357.
 13. Murthy J (2010) Neurological complications of dengue infection. *Neurol India* 58: 581-584.
 14. Parkash O, Almas A, Jafri Wasmin SM, Hamid S, Akhtar J, Alishah H (2010) Severity of acute hepatitis and its outcome in patients with dengue fever in a tertiary care hospital Karachi, Pakistan (South Asia). *BMC Gastroenterology* 10: 43.
 15. Itha S, Kashyap R, Krishnani N, Saraswat VA, Choudhri G, Aggarwal R (2005) Profile of liver involvement in dengue virus infection. *Natl Med J India* 18: 127-130.
 16. de Souza LJ, Goncalves Carneiro H, Souto Filho JT, Ferreira de Souza T, Azevedo Cortes V, Neto CG, Bastos Assed D, Wallace da Silva Siqueira E (2002) Hepatitis in dengue shock syndrome. *Braz J Infect Dis* 6: 322-327.
 17. Kumar R, Tripathi P, Tripathi S, Kanodia A, Venkatesh V (2008) Prevalence of dengue infection in north Indian children with acute hepatic failure. *Ann Hepatol* 7: 59-62.
 18. Huang YH, Liu CC, Wang ST, Lei HY, Liu HS, Lin WS, Wu HL, Yeh TM (2001) Activation of coagulation and fibrinolysis during dengue virus infection. *Med Virol* 63: 247-251.
 19. Kumar R, Tripathi S, Tambe JJ, Arora V, Nag VL (2008) Dengue encephalopathy in children in Northern India: Clinical features and comparison with non dengue. *J Neurol Sci* 269: 41-48.

Corresponding author

Ritu Karoli
Department of Medicine
Era's Lucknow Medical College
Sarfarazganj, Hardoi Road
Lucknow, Uttar Pradesh, India

Conflict of interests: No conflict of interests is declared.