

Original Article

## Relationship between age and intensive care unit-acquired bloodstream infections in infectious disease patients in Croatia

Mirjana Balen Topić<sup>1</sup>, Marija Santini<sup>1</sup>, Bruno Baršić<sup>1</sup>

<sup>1</sup> School of Medicine, University of Zagreb, University Hospital for Infectious Diseases "Dr. Fran Mihaljević", Zagreb, Croatia

### Abstract

**Introduction:** Intensive care unit-acquired bloodstream infections (ICU-BSI) belong to the most important nosocomial infections. Since there is scarce data available on their relationship with older age, we performed this study to estimate the age-related incidence of ICU-BSI and the odds of acquiring ICU-BSI in elderly critically ill infectious disease patients.

**Methodology:** A retrospective observational analysis of prospectively collected demographic and clinical data of adult mechanically ventilated infectious disease patients, treated in a teaching hospital in Croatia between 1994 and 2008, using univariate, bivariate, and multivariate logistic regression analyses.

**Results:** Of the 1,093 included patients, 509 (46.6%) were  $\geq 65$  years old, among 256 (23.4%) of whom a total of 353 ICU-BSI episodes were recorded. No significant difference among ICU-BSI causative microorganisms between the observed age groups was found ( $P = 0.4940$ ). The rate of patients with ICU-BSI was higher among elderly ones (26.1 vs. 21.1%,  $P = 0.048$ ), and elderly patients used the ICU facilities (ICU stay, duration of mechanical ventilation and central venous catheter [CVC] use) significantly longer ( $P < 0.05$ ). However, older age was not positively related with the development of ICU-BSI (OR 0.99, 95% CI: 0.71-1.38); as opposed to the duration of CVC use (OR 1.09, 95% CI: 1.07-1.10).

**Conclusion:** It seems that among adult mechanically ventilated infectious disease patients, borderline significantly higher rate of ICU-BSI among those aged  $\geq 65$  years was related to longer use of ICU facilities, rather than to their older age itself. The duration of CVC use was identified as the only factor positively related to the development of ICU-BSI.

**Key words:** elderly; intensive care unit; bloodstream infections.

*J Infect Dev Ctries* 2018; 12(5):352-358. doi:10.3855/jidc.9950

(Received 17 November 2017 – Accepted 09 February 2018)

Copyright © 2018 Balen Topić *et al.* This is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Introduction

Population aging is a well-recognized phenomenon globally, and elderly people have become more frequent users of medical care services in general. According to previous reports from different countries, the proportion of ICU-treated elderly patients ranges between 26-52% [1,2] and a shift in primary ICU diagnoses from cardiovascular to infectious diseases in elderly patients has been observed in a recent study [3]. In comparison with the younger population, community-acquired infections are more common and associated with a poorer outcome in elderly patients [4]. The reasons for this increased susceptibility include epidemiological elements, immunosenescence, malnutrition as well as a large number of age-associated physiological and anatomical alterations [5].

The impact of ageing on the incidence of nosocomial infections seems to be a controversial issue. Keeping in mind the differences in study designs, used statistical methods, sample sizes and patients' medical

features, which could influence conclusions, when we observed studies that compared population aged  $\geq 65$  years with younger adults, some studies found higher incidence of nosocomial, particularly ICU-acquired infections among elderly patients, [6-8] while other failed to do so [9-12]. Despite of the assumed increased susceptibility for infection, few studies have shown that the incidence of ICU-BSI even decreased with age [13,14].

ICU-BSI cases are considered as the most common, most lethal and most expensive nosocomial infections, [15-17] reaching in elderly patients in developed countries mean attributable costs of \$43,208 per episode [18]. It has been estimated that ICU-BSI complicate between 4.4 to 6.8% of admissions lasting longer than 48 to 72 hours [13,15,19].

Croatian population is regarded as one of the oldest in Europe. We performed an observational study in a large cohort of mechanically ventilated, critically ill

infectious disease patients, with the aim to estimate the age-related incidence and odds for acquiring ICU-BSI.

## Methodology

### *Study of population and data sources*

We performed a retrospective observational analysis of prospectively collected data of age-related ICU-BSI incidence among critically ill, mechanically ventilated infectious disease patients, treated at the 18-bedded adult ICU of the University Hospital for Infectious Diseases in Zagreb, Croatia, in the period from 1994 to 2008. Data were obtained from electronic database of ICU admissions, and hospital microbiology records of positive blood cultures. Standard protocols for the collection, analysis, and reporting of blood cultures were employed. For the analysis of routinely collected data the study was approved by the institutional review boards of the study hospital as well as the affiliated School of Medicine, waiving the requirement for specific patient consent.

### *Definitions*

We used the American College of Chest Physicians/Society of Critical Care Medicine (ACCP/SCCM) definitions of ICU-BSI; and considered both primary and secondary BSIs in our analysis [20]. ICU-BSI was defined as the presence of clinical evidence of infection, accompanied by positive blood cultures for a bacterium or fungus obtained at the same time, but more than 48 hours after the admission to the ICU. In accordance with the US Centers for Disease Control and Prevention guidelines, we did not include cultures of coagulase-negative staphylococci or other common commensal skin organisms, unless two cultures separately isolated the same species of microorganism [21]. To allow the study of a population at risk, we excluded all ICU admissions lasting less than 48 hours. Elderly patients were defined as equal to or older than 65 years. The results were compared to patients aged 18-64 years.

### *Data analysis*

The following data were analyzed: patients' demographics, previous health status (according to Knaus classification), Glasgow Coma Score (GCS) on admission, Acute Physiology and Chronic Health Evaluation II (APACHE II) physiology score on admission, APACHE II score minus age points, type of primary infection on admission regarding central nervous system (CNS) involvement, usage of CVC, duration of CVC usage (in days), duration of mechanical ventilation (MV) (in days), incidence of

ICU-BSI, causative microorganism of ICU-BSI, incidence of other nosocomial infections (pneumonia, urinary tract infections), and duration of ICU stay. As an infection with CNS involvement we also considered sepsis with secondary CNS infection and tetanus.

Univariate analyses of continuous variables were reported as a mean value and standard deviation, and categorical variables as frequency and percentage. For comparison of the groups for continuous variables, the Mann-Whitney U test was used, and for categorical variables, the chi-squared test or Fisher's exact test, when appropriate. We described an association of predictors with the dependent variable being the development of ICU-BSI with an "odds ratio" (OR). Logistic regression multivariate analysis was performed to test whether increased age is an independent factor for ICU-BSI development after adjustment for possible confounding factors. Besides age group, all variables which in univariate analysis showed significant association with BSI were included in the model. The Hosmer-Lemeshow test was used as a goodness-of-fit statistics. Statistical significance was defined as  $P < 0.05$ . Statistical analysis was performed using SAS 9.2 software, SAS Institute Inc., 2009, Cary, NC, USA.

## Results

Of the 2,171 patients with severe infections treated in the ICU, 922 were excluded from the study due to lack of mechanical ventilation, and further 156 because of ICU-stay shorter than 48h. Among the remaining 1,093 included patients, 509 (46.6%) were  $\geq 65$  years (elderly group), while 584 (53.4%) were younger adults. In 256 (23.4%) observed patients a total of 353 ICU-BSI episodes were recorded. Among them, 195 episodes of ICU-BSI were recorded in 26.1% (133/509) elderly patients, and 158 episodes in 21.1% (123/584) younger adult patients.

The most common causative microorganisms were non-fermenting gram-negative bacteria causing 32.0% of a total of 353 ICU-BSI episodes, followed by *Enterobacteriaceae* (sensitive 9.9%, and 3<sup>rd</sup> generation cephalosporins resistant/ESBL 14.2%), *Staphylococcus aureus* (MRSA 9.3% and MSSA 2.0%), *Candida sp.* (7.1%), *Enterococcus sp.* (2.9% ampicillin sensitive and 1.1% ampicillin resistant), coagulase-negative staphylococcus (2.8%) and other bacteria (1.1%), respectively; however, 17.6% of ICU-BSIs were polymicrobial. No significant difference in the distribution of ICU-BSI causative microorganisms between the two observed age groups was found ( $p = 0.494$ ).

**Table 1.** The comparison of basic demographic and clinical data and outcomes between two age groups of mechanically ventilated, critically ill infectious disease patients.

Variables	Patient age group (N = 1093)		P
	18-64 years (N = 584)	≥ 65 years (N = 509)	
Age mean (SD)*	47.6 (12.8)	74.9 (6.5)	
Female N (%)	211 (36.1)	259 (50.9)	< 0.001
Male N (%)	373 (63.9)	250 (49.1)	
Knaus classification N (%)			0.005
Healthy	213 (36.5)	141 (27.7)	
Mild chronic disease	215 (36.8)	224 (44.0)	
Severe chronic disease	126 (21.6)	105 (20.6)	
Terminal chronic disease	30 (5.1)	39 (7.7)	
Infection with CNS <sup>§</sup> involvement N (%)	321 (55.0)	217 (42.6)	< 0.001
Infection without CNS involvement N (%)	263 (45.0)	292 (57.4)	
Mean APACHE II <sup>1</sup> (SD)	19.1 (8.1)	22.3 (8.8)	< 0.001
Mean age-adduced APACHE II (SD)	17. (7.9)	16.9 (8.7)	0.106
Mean GCS <sup>2</sup> (SD)	10.1 (4.3)	10.9 (4.1)	0.001
CVC <sup>3</sup> use N (%)	549 (94.0)	484 (95.1)	0.437
Mean duration of CVC use in days (SD)	13.1 (12.2)	15.8 (14.6)	0.005
Mean duration of MV <sup>4</sup> in days (SD)	12.7 (14.1)	16.2 (20.4)	0.005
Mean duration of ICU stay in days (SD)	18.5 (18.1)	22.9 (26.4)	0.005
Episodes of nosocomial UTI <sup>5</sup> (% of patients)	94 (14.4)	141 (22.4)	0.005
Episodes of nosocomial pneumonia (% of patients)	90 (13.9)	72 (13.2)	0.917
Patients with ICU-BSI <sup>6</sup> N (%)	123 (21.1)	133 (26.1)	0.048
Incidence of ICU-BSI/1,000 CVC days	16.4	17.7	0.046
Incidence of ICU-BSI/1,000 ICU days	12.3	12.4	0.097
Overall mortality N (%)	250 (42.8)	268 (52.8)	< 0.001

\*SD: Standard deviation; <sup>§</sup>CNS: Central Nervous System; <sup>1</sup>APACHE II: Acute Physiology and Chronic Health Evaluation score II; <sup>2</sup>GCS: Glasgow Coma Score; <sup>3</sup>CVC: Central Venous Catheter; <sup>4</sup>MV: Mechanical Ventilation; <sup>5</sup>UTI: Urinary Tract Infection; <sup>6</sup>ICU-BSI: Intensive- Care-Unit-acquired Bloodstream Infection.

**Table 2.** The comparison of basic demographic and clinical data between groups with and without intensive care unit-acquired bloodstream infection among mechanically ventilated, critically ill infectious disease patients.

Variables	ICU-BSI* (N = 1093)		P
	No (N = 837)	Yes (N = 256)	
Age mean (SD) <sup>§</sup>	59.5 (17.6)	62.9 (15.1)	0.017
Age group N (%)			0.048
18-64 years	461 (55.1)	123 (48.0)	
≥ 65 years	376 (44.9)	133 (52.0)	
Sex N (%)			< 0.001
Female	334 (39.9)	136 (53.1)	
Male	503 (60.1)	120 (46.9)	
Knaus classification N (%)			0.083
Healthy	276 (33.0)	78 (30.5)	
Mild chronic disease	320 (38.2)	119 (46.5)	
Severe chronic disease	188 (22.5)	43 (16.8)	
Terminal chronic disease	53 (3)	16 (6.3)	
Infection with CNS <sup>1</sup> involvement N (%)			< 0.001
Yes	386 (46.1)	152 (59.4)	
No	451 (53.9)	104 (40.6)	
APACHE II <sup>2</sup> mean (SD)	21.1 (8.6)	19.0 (8.1)	< 0.001
Age-adduced APACHE II mean (SD)	17.7 (8.3)	15.3 (8.1)	< 0.001
GCS <sup>3</sup> mean (SD)	10.2 (4.2)	11.2 (4.2)	< 0.001
Duration of CVC <sup>4</sup> use in days mean (SD)	10.9 (9.6)	25.6 (17.4)	< 0.001
Duration of MV <sup>5</sup> in days mean (SD)	10.1 (10.3)	28.3 (26.4)	< 0.001
Duration of ICU <sup>6</sup> stay in days mean (SD)	15.1 (14.3)	38.4 (32.8)	< 0.001
Overall mortality N (%)	416 (49.7)	103 (40.2)	0.008

\*ICU-BSI: Intensive Care Unit-acquired Bloodstream Infection; <sup>§</sup>SD: Standard deviation; <sup>1</sup>CNS: Central Nervous System; <sup>2</sup>APACHE II: Acute Physiology and Chronic Health Evaluation score II; <sup>3</sup>GCS: Glasgow Coma Score; <sup>4</sup>CVC: Central Venous Catheter; <sup>5</sup>MV: Mechanical Ventilation; <sup>6</sup>ICU: Intensive Care Unit.

Among the two observed age groups, a significantly higher mean duration of CVC use, duration of MV and ICU stay, as well as higher mortality rate was found in the elderly, while the rate of patients with ICU-BSI, and the incidence of ICU-BSI per 1,000 CVC days were borderline significantly higher in elderly patients; detailed comparison of baseline demographic and clinical data and outcomes is shown in Table 1.

Bivariate analyses of groups with and without ICU-BSI, according to demographic and clinical data, and ICU facility use, shown in Table 2, identified the following variables ( $P < 0.05$ ) as variables that could influence the development of ICU-BSI: sex, age group, type of primary infection regarding involvement of CNS, APACHE II score minus age points, duration of MV as well as the duration of CVC use. These variables (except the duration of MV, because of its interaction with duration of CVC use) were included in logistic regression analyses as independent variables. The results of multivariate analysis are shown in Table 3. The model fitted well showing a high explanatory value ( $c = 0.802$ ). The incidence of ICU-BSI was equally distributed between two age groups (OR: 0.99, 95% CI: 0.71-1.38). APACHE II score corrected for age (OR 0.97, 95% CI: 0.95-0.99) and the overall duration of CVC use (OR 1.09, 95% CI: 1.07-1.10) were two parameters independently associated with ICU-BSI.

## Discussion

With the average age of 41,7 years and with 17,7% of the ~4,285 million population being  $\geq 65$  years old in 2011, most likely due to the low natality and high emigration rate among young people, Croatian population is one of the five oldest in Europe [22]. Almost half of our ICU patients satisfy the criterion of being elderly, and we expected the greater risk of ICU-BSI in this group of patients. Contrary to our

expectations, the results of this study show equal odds for the development of ICU-BSI among adult, critically ill infectious disease patients  $< 65$  and those  $\geq 65$  years.

A negative relation between increasing age and the risk of ICU-BSI development, and incidence of ICU-BSI, respectively, has been found in two large retrospective studies (Prowle *et al.*, 2011, Blot *et al.*, 2009.) [13,14]. These unexpected results were explained with assumed under-diagnosing of ICU-BSI among the elderly, because of the association of older age with a weaker inflammatory response that could decrease the likelihood of blood culture sampling and reduce the observed incidence of BSI in older patients [5].

Microbiological characteristics of the ICU environment, ICU procedures, or even the impact of severe primary disease on defense mechanisms of ICU patients have been suggested as factors that affect the development of ICU-acquired infections.

A positive relationship between the duration of CVC use and the risk of developing a CVC-related and primary ICU-BSI has been well documented [23,24], but it seems that the incidence of CVC-related ICU-BSI is 3-5 times higher in developing countries in comparison to developed countries [25]. According to Maki's observations, 2/3 of primary nosocomial BSIs also originate from contaminated CVC [26]. The high rate of 81% of CVC-related BSI has been observed, and the presence of CVC was identified as an independent risk factor for nosocomial BSI development among elderly patients (OR 7.5, 95% CI: 2.5-22.9) [18,27]. The unusually high incidence of ICU-BSI expressed per 1,000 CVC days in our patients (16.4 in younger adults and 17.7 in elderly) is due to the jointly calculated incidence of CVC-related, primary and secondary ICU-BSIs in our study, and those incidences are approximately three times higher in comparison to

**Table 3.** Analysis of the correlation between significant factors and the development of intensive care unit-acquired bloodstream infection using logistic regression.

Variables	Probability of ICU-BSI* development		
	Odds Ratio	95% Confidence intervals (Wald interval)	
Age (years): 18-64 / $\geq 65$	0.99	0.71	1.38
Sex: female / male	1.29	0.93	1.79
CNS <sup>§</sup> infections / other infections	1.17	0.83	1.63
APACHE II <sup>1</sup> minus age points	0.97	0.95	0.99
Duration of CVC <sup>2</sup> use (days)	1.09	1.07	1.10

\*ICU-BSI: Intensive-Care-Unit-acquired Bloodstream Infection; <sup>§</sup>CNS: Central Nervous System; <sup>1</sup>APACHE II: Acute Physiology and Chronic Health Evaluation score II; <sup>2</sup>CVC: Central Venous Catheter.



those reported solely for CVC-related ICU-BSIs [23,28,29]. The multivariate analysis results have shown that the duration of CVC use was the only analyzed factor positively related to the development of ICU-BSI among our patients, which highlights the importance of CVC use in the epidemiology and pathogenesis of ICU-BSI. However, a significant decrease in the incidence of CVC-related ICU-BSIs from 5.7 to 1.5 per 1,000 CVC days, after the implementation of preventive measures, has been reported from Brasil [30], and a sustained very low incidence of CVC-related ICU-BSIs of 0.6/1,000 CVC days from a highly developed country in a large Swedish prospective study [31].

As expected, our two observed age groups differed significantly according to previous health status assessed by the Knaus classification, with higher rate of elderly patients belonging to the group with mild chronic disease (44.0 vs. 36.8%) and with terminal chronic disease (7.7 vs. 5.1%), in comparison to younger adults; however, the difference of mean age-adjusted APACHE II scores at admission between the observed age groups was not found.

Conversely to our results, many previous studies have shown that greater illness severity on ICU-admission, expressed with APACHE II or APACHE III score, is related to the higher risk of developing ICU-BSI [13,32-34]. The finding of a negative relationship between APACHE II minus age points at ICU-admission, and the development of ICU-BSI in our study, suggests a high rate of early mortality among critically ill patients with domicile infections, which has lowered their chance for development of ICU-BSI.

Due to longer use of ICU facilities (CVC, MV and ICU stay) in our patients, the observed overall ICU-BSI rate of 26.1% in elderly and 21.1% in younger adults were much higher in comparison to those reported from other studies, which ranged between 2.7-9.0% [15,19,33,35-37]. Relatively expressed ICU-BSI incidence of 12.4 ICU-BSIs in elderly and 12.3 in younger adults per 1,000 ICU days in our observed severe infectious disease patients fits among the results of other studies, where that incidence in patients treated at ICU, because of different medical and surgical conditions, was mostly in the range of 5-19/1,000 ICU days [19,35,37]. Exceptionally, the highest recorded incidence, reaching 44/1,000 ICU days, has been reported from Upper Egypt [38]. However, unlike data collected by the Turkish authors who reported a significantly higher incidence of ICU-BSIs among elderly patients in comparison to younger adults (17.37

vs. 11.21/1,000 ICU days), in our study age-based difference was not found [8].

The main shortcomings of this study are the disadvantages associated with its retrospective design. Also, although the signs of infection were carefully monitored in our critically ill infectious disease patients, the possibility of underdiagnosing ICU-acquired infections in elderly patients cannot be completely excluded.

## Conclusion

It seems that among adult mechanically ventilated infectious disease patients, borderline significantly higher rate of ICU-BSI among those aged  $\geq 65$  years was related to longer use of ICU facilities, rather than to their older age itself. The duration of CVC use was identified as the only factor positively related to the development of ICU-BSI.

## Funding

Data used for this study were collected within the framework of Research Project No. 108-10800002-0102, which was partially funded by a grant from the Ministry of Science, Education and Sports of the Republic of Croatia.

## References

1. Hennessy D, Juzwishin K, Yergens D, Noseworthy T, Doig C (2005) Outcomes of elderly survivors of intensive care: A review of the literature. *Chest* 127: 1764-1774.
2. Sodhi K, Singla MK, Shrivastava A, Bansal N (2014) Do Intensive Care Unit treatment modalities predict mortality in geriatric patients: An observational study from an Indian Intensive Care Unit. *Indian J Crit Care Med* 18: 789-795.
3. Sjoding MW, Prescott HC, Wunsch H, Iwashyna TJ, Cooke CR (2016) Longitudinal Changes in ICU Admissions Among Elderly Patients in the United States. *Crit Care Med* 44: 1353-1360.
4. Yoshikawa TT (2000) Epidemiology and unique aspects of aging and infectious diseases. *Clin Infect Dis* 30: 931-933.
5. Gavazzi G, Krause KH. Ageing and infection (2002) *Lancet Infect Dis* 2: 659-666.
6. Solis-Hernandez PS, Vidales-Reyes M, Garza-Gonzalez E, Guajardo-Alvarez G, Chavez-Moreno S, Camacho-Ortiz A (2016) Hospital-acquired infections in elderly versus younger patients in an acute care hospital. *Int J Infect* 3: e32620.
7. Avici M, Ozgenc O, Coskuner SA, Olut AI (2012) Hospital acquired infections (HAI) in the elderly: comparison with the younger patients. *Arch Gerontol Geriatr* 54: 247-250.
8. Batirel, A., Benzonana, N., Ak, O., Kavak, M., Akay, D., Cag, Y., & Ozer, S. (2014). Hospital-acquired infections in the elderly in intensive care unit. In J. Vila (Ed.), *Proceedings of 24th European Congress of Clinical Microbiology and Infectious Diseases* (P1606). Barcelona, Spain: European Society of Clinical Microbiology and Infectious Diseases..
9. Tang EY, Hsu LF, Lam KN, Pang WS (2003) Critically ill elderly who require mechanical ventilation: The effects of age

- on survival outcomes and resource utilisation in the medical intensive care unit of a general hospital. *Ann Acad Med Singapore* 32: 691-696.
10. Erbay H, Yalcin AN, Serin S, Turgut H, Tomatir E, Cetin B, Zencir M (2003) Nosocomial infections in intensive care unit in a Turkish university hospital: A 2-year survey. *Intensive Care Med* 29: 1482-1488.
  11. Appelgren P, Hellstrom I, Weitzberg E, Soderlund V, Bindslev L, Ransjo U (2001) Risk factors for nosocomial intensive care infection: a long-term prospective analysis. *Acta Anaesthesiol Scand* 45: 710-719.
  12. Sacanella E, Pérez-Castejón JM, Nicolás JM, Masanés F, Navarro M, Castro P, López-Soto A (2009) Mortality in healthy elderly patients after ICU admission. *Intensive Care Med* 35: 550-555.
  13. Prowle JR, Echeverri JE, Ligabo EV, Sherry N, Taori GC, Crozier TM, Hart GK, Korman TM, Mayall BC, Johnson PD, Bellomo R (2011) Acquired bloodstream infection in the intensive care unit: incidence and attributable mortality. *Crit Care* 15: R100.
  14. Blot S, Cankurtaran M, Petrovic M, Vandijck D, Lizy C, Decruyenaere J, Danneels C, Vandewoude K, Piette A, Vershraegen G, Van Den Noortgate N, Peleman R, Vogelaers D (2009) Epidemiology and outcome of nosocomial bloodstream infection in elderly critically ill patients: A comparison between middle-aged, old, and very old patients. *Crit Care Med* 37: 1634-1641.
  15. Garrouste-Orgeas M, Timsit JF, Tafflet M, Misset B, Zahar JR, Soufir L, Lazard T, Jamali S, Mourvillier B, Cohen Y, De Lasseuse A, Azoulay E, Cheval C, Descorps-Declare A, Adrie C, Costa de Beauregard MA, Carlet J; OUTCOMES Study Group (2006) Excess risk of death from intensive care unit-acquired nosocomial bloodstream infections: A reappraisal. *Clin Infect Dis* 42: 1118-1126.
  16. Wisplinghoff H, Bischoff T, Tallent SM, Seifert H, Wenzel RP, Edmond MB (2004) Nosocomial bloodstream infections in US hospitals: analysis of 24,179 cases from a prospective nationwide surveillance study. *Clin Infect Dis* 39: 309-317.
  17. Laupland KB, Lee H, Gregson DB, Manns BJ (2006) Cost of intensive care unit-acquired bloodstream infections. *J Hosp Infect* 63: 124-32.
  18. Kaye KS, Marchaim D, Chen TY, Baures T, Anderson DJ, Choi Y, Sloane R, Schmader KE (2014) Effect of nosocomial bloodstream infections on mortality, length of stay, and hospital costs in older adults. *J Am Geriatr Soc* 62: 306-311.
  19. Laupland KB, Zygun DA, Davies HD, Church DL, Louie TJ, Doig CJ (2002) Population-based assessment of intensive care unit-acquired bloodstream infections in adults: Incidence, risk factors, and associated mortality rate. *Crit Care Med* 30: 2462-2467.
  20. Bone RC, Balk RA, Cerra FB, Dellinger RP, Fein AM, Knaus WA, Schein RM, Sibbald WJ (1992) Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. *Chest* 101: 1644-1655.
  21. Horan TC, Andrus M, Dudeck MA (2008) CDC/NHSN surveillance definition of health care-associated infection and criteria for specific types of infections in the acute care setting. *Am J Infect Control* 36: 309-332.
  22. Lj Ostroski (2013) 1468/2012 Census of population, households and dwellings, 2011, population by sex and age: statistical reports. Zagreb, Croatia: Croatian Bureau of Statistics. Available [https://www.dzs.hr/Hrv\\_Eng/publication/2012/SI-1468.pdf](https://www.dzs.hr/Hrv_Eng/publication/2012/SI-1468.pdf). Accessed 29 May 2018.
  23. van der Kooij TI, de Boer AS, Manniën J, Wille JC, Beaumont MT, Mooi BW, van den Hof S (2007) Incidence and risk factors of device-associated infections and associated mortality at the intensive care unit in the Dutch surveillance system. *Intensive Care Med* 33: 271-278.
  24. Rojo D, Pinedo A, Clavijo E, García-Rodríguez A, García V (1999) Analysis of risk factors associated with nosocomial bacteraemias. *J Hosp Infect* 42: 135-141.
  25. Rosenthal VD, Maki DG, Jamulitrat S, Medeiros EA, Todi SK, Gomez DY, Leblebicioglu H, Abu Khader I, Miranda Novales MG, Berba R, Ramirez Wong FM, Barkat A, Pino OR, Dueñas L, Mitrev Z, Bijie H, Gurskis V, Kanj SS, Mapp T, Hidalgo RF, Ben Jaballah N, Raka L, Gikas A, Ahmed A, Thu le TA, Guzmán Sirtt ME (2010) International Nosocomial Infection Control Consortium (INICC) report, data summary for 2003-2008, issued June 2009. *Am J Infect Control* 38: 95-106.
  26. Maki DG (1981) Nosocomial bacteremia: an epidemiologic overview. *Am J Med* 70: 719-732.
  27. Reunes S, Rombaut V, Vogelaers D, Brusselaers N, Lizy C, Cankurtaran M, Labeau S, Petrovic M, Blot S (2011) Risk factors and mortality for nosocomial bloodstream infections in elderly patients. *Eur J Intern Med* 22: e39-44.
  28. McLaws ML, Taylor PC (2003) The Hospital Infection Standardised Surveillance (HISS) programme: analysis of a two-year pilot. *J Hospit Infect* 53: 259-267.
  29. Napalkov P, Felici DM, Chu LK, Jacobs JR, Begelman SM (2013) Incidence of catheter-related complications in patients with central venous or hemodialysis catheters: a health care claims database analysis. *BMC Cardiovasc Disord* 13: 86.
  30. Marra AR, Cal RGR, Durão MS, Correa L, Guastelli LR, Moura DF Jr, Edmond MB, Dos Santos OF (2010) Impact of a program to prevent central-line-associated bloodstream infection in the zero tolerance era. *Am J Infect Control* 38: 434-439.
  31. Hammarskjöld F, Berg S, Hanberger H, Taxbro K, Malmvall BE (2014) Sustained low incidence of central venous catheter-related infections over six years in a Swedish hospital with an active central venous catheter team. *Am J Infect Control* 42: 122-128.
  32. Laupland KB, Gregson DB, Zygun DA, Doig CJ, Mortis G, Church DL (2004) Severe bloodstream infections: a population-based assessment. *Crit Care Med* 32: 992-997.
  33. Magnason S, Kristinsson KG, Stefansson T, Erlendsdottir H, Jonsdottir K, Kristjansson M, Jonmundsson E, Baldursdottir L, Sigvaldason H, Gudmundsson S (2008) Risk factors and outcome in ICU-acquired infections. *Acta Anaesthesiol Scand* 52: 1238-1245.
  34. Pratikaki M, Platsouka E, Sotiropoulou C, Vassilakopoulos T, Paniara O, Roussos C, Routsis C (2009) Risk factors for and influence of bloodstream infections on mortality: a 1-year prospective study in a Greek intensive-care unit. *Epidemiol Infect* 137: 727-735.
  35. Pittet D, Tarara D, Wenzel RP (1997) Nosocomial bloodstream infection in critically ill patients: excess length of stay, extra costs, and attributable mortality. *JAMA* 271: 1598-1601.
  36. Laupland KB, Kirkpatrick AW, Church DL, Ross T, Gregson DB (2004) Intensive-care-unit-acquired bloodstream infections in a regional critically ill population. *J Hosp Infect* 58: 137-145.

37. Pittet D, Thiévent B, Wenzel RP, Li N, Auckenthaler R, Suter PM (1996) Bedside prediction of mortality from bacteremic sepsis: a dynamic analysis of ICU patients. *Am J Respir Crit Care Med* 153: 684-693.
38. Ahmed SH, Daef EA, Badary MS, Mahmoud MA, Abd-Elsayed AA (2009) Nosocomial bloodstream infection in intensive care units at Assuit University Hospitals (Upper Egypt) with special reference to extended spectrum  $\beta$ -lactamase producing organisms. *BMC Res Notes* 2: 76.

**Corresponding author**

Assist Prof Mirjana Balen Topić, MD, PhD  
University Hospital for Infectious Diseases “Dr. Fran Mihaljević”  
Mirogojska 8, 10 000 Zagreb, Croatia  
Phone: ++385 91 4012 537  
Fax: ++385 1 2826 153  
Email: mbalen@bfm.hr

**Conflict of interests:** No conflict of interests is declared.